

Near Extinction of a Large, Widely Distributed Fish

Jill M. Casey and Ransom A. Myers*

Are extinctions of marine vertebrates as rare and unlikely as current data indicate? Long-term research surveys on the continental shelf between the Grand Banks of Newfoundland and southern New England reveal that one of the largest skates in the northwest Atlantic, the barndoor skate (*Raja laevis*), is close to extinction. Forty-five years ago, research surveys on St. Pierre Bank (off southern Newfoundland) recorded barndoor skates in 10% of their tows; in the last 20 years, none has been caught, and this pattern of decline is similar throughout the range of the species.

Elasmobranchs, such as sharks and rays, tend to be very susceptible to the effects of fishing because they generally grow slowly, mature late in life, and produce few offspring (1). Species of the family Rajidae, although the most fecund of the elasmobranchs (2), are known to experience varying degrees of resilience to exploitation (3) because of the large range in life history characteristics within this family. Although shark fisheries usually cause a sharp decline in species abundance (4), dogfish and skates on Georges Bank have increased in biomass after the depletion of groundfish stocks (5). It has been suggested that the energy released into the ecosystem by these depleted stocks has provided resources for elasmobranch populations to increase. The recent introduction of a directed fishery for dogfish and skate on Georges Bank, however, has resulted in a marked decline of these species (6).

The barndoor skate, *Raja laevis*, is one of the largest skates in the northwest Atlantic (7), and it ranges from Cape Hatteras to the Grand Banks of Newfoundland (Fig. 1). Once common (8), this distinctive species, with a maximum body width of just over 1 m, now appears to be near extinction. Although the extinction of marine species is thought to be rare (9), the closely related "common" skate in the northeast Atlantic, *Raja batis*, was shown to be locally extinct in the Irish Sea

(10). If current population trends continue, however, the barndoor skate could become the first well-documented example of extinction in a marine fish species.

Biomass (in kilograms per square kilometer) of the barndoor skate (Fig. 2) was determined from research vessel surveys (11) that have been conducted in the spring on the southern Grand Bank and St. Pierre Bank since 1951, in the summer on the Scotian Shelf since 1970, and in the autumn from Georges Bank to southern New England since 1963. The population trend is similar for all regions, with biomass decreasing into the early 1970s, after which barndoor skates were caught only on Browns Bank and nearby Georges Bank (12).

The longest time series available are for the southern Grand Bank, at the species' northern limit on the continental shelf, and St. Pierre Bank, where barndoor skates were

once commonly found. Compared with other skate species on St. Pierre Bank, the barndoor skate had been one of the most numerous skates, second in abundance only to the thorny skate (*Raja radiata*) (13). If we consider the mean biomass of the barndoor skate in each decade and the corresponding mean weight of individuals on St. Pierre Bank, the average number of barndoor skates in the 1950s would have been on the order of 0.6 million. That number would decrease to about 0.2 million individuals in the 1960s and to less than an estimated 500 individuals in the 1970s. The other smaller skate species, namely thorny skate and smooth skate (*Raja senta*), were actually increasing in biomass over this time period.

The mean weight of individual skates on St. Pierre Bank is a good indicator of their vulnerability to commercial trawls. Species large enough to be caught in trawls have been decreasing in size over the time period, whereas the size of the smallest species, the smooth skate, has remained fairly constant. The regulated mesh size on commercial fishing gear has ranged from 7 to 14 cm over the survey period, although a smaller mesh size has been used illegally (14). The barndoor skate is vulnerable to commercial trawls from hatching at a length of about 20 cm (7).


Direct biological information on skates in the northwest Atlantic is scarce. Sufficient comparative information, however, is available to estimate the mortality required to drive this species to extinction. The closest relative of the barndoor skate in the North Atlantic, the common skate (7), matures at about 11 years (10). We should expect a similar age at maturity on Georges Bank, which has a similar temperature regime (15). Maximum egg production, which can be estimated from the inverse relation with the weight of the young at hatching (16), is about 47 eggs per year.

Considering the age at maturity and the annual fecundity of the barndoor skate, the

J. M. Casey, Department of Biology, Memorial University of Newfoundland, St. John's, Newfoundland, Canada A1C 5S7. R. A. Myers, Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada B3H 4J1.

*To whom correspondence should be addressed. E-mail: ransom.myers@dal.ca

Fig. 1. Map of the Northwest Atlantic Fisheries Organization. Subdivisions in which populations were assessed in this analysis are shown. The 300-m isobath (dotted line) is given for reference. 3N and 3O, southern Grand Bank; 3Ps, St. Pierre Bank; 4Vn, Sydney Bight; 4Vs, Banquereau Bank; 4W, Sable Island Bank; 4X, Browns Bank; 5Y, Gulf of Maine; 5Ze, Georges Bank; 5Zw, southern New England. The numbers on the axes are degrees of latitude and longitude.



instantaneous mortality rate required to drive this species to extinction is about 0.4 per year, assuming that mature and immature mortalities are equal (17). At the northern limit of the barndoor skate's range, where the bottom temperature averages 2.5°C [about 6°C colder than on Georges Bank (15)], the age at maturity should be about double (18), and the instantaneous mortality required to drive the species to extinction would be expected to be closer to 0.2.

The fishing mortalities for cod (*Gadus morhua*) have been above 0.4 and often much greater in these regions for more than 30 years (19). It is therefore not surprising that the barndoor skate population would be reduced to such extreme levels when taken

as by-catch in major fisheries, for example, cod and redfish (*Sebastes* sp.). For a species that matures at such a late age (at least 11 years), there is a greater probability that individuals will be caught before they are able to reproduce. This situation is amplified for populations at the northern limit of the range.

The only recent research survey catches of barndoor skate in the Newfoundland area have been very deep (greater than 1000 m) and in an area north of the reported range of this species, where surveys were conducted in support of a new fishery for Greenland halibut (*Reinhardtius hippoglossoides*). Whether or not barndoor skates were commonly found at this depth over

the entire range is not known, as research surveys were rarely conducted at such depths. These great depths, however, may have represented one of the last refuges for this species.

Barndoor skates have survived on Browns Bank and Georges Bank probably because of a faster growth rate combined with the seasonal closure of parts of these banks to trawling (20). Perhaps the only hope for the long-term survival of this species is to designate an area protected from trawling on all the banks that is sufficiently large to allow for a self-sustaining population. A protected area would also provide a simple and effective means to conserve other species.

Failure to examine historical data has resulted in the largest skate in the northwest Atlantic being driven to near extinction without anyone noticing. If such a large, easily identified species has been allowed to disappear in an area that is well surveyed, the fate of little known species is likely to be worse. This study shows the importance of assembling data on as long a time scale as possible and as wide a spatial scale as possible; otherwise, the near extinction of a very apparent species may be missed.

A directed fishery for skate was initiated off the coasts of Newfoundland and Nova Scotia in 1994 as an attempt to use nontraditional groundfish species. Elasmobranchs experience low natural mortality when compared with teleost fish species and, at low population levels, are not able to compensate, for example, with increased fecundity or reduced natural mortality (4). Consequently, the end result of increased mortality due to fishing pressure could be extinction.




Fig. 2. Estimates of absolute biomass for barndoor skate (*R. laevis*) from the southern Grand Bank (the northern limit of the range) to southern New England (close to the southern limit of the range). Open circles are zero catches. An exponential decay curve ($Ne^{-\delta t}$) was fit to the data with nonlinear least squares, where N is the population size in the first year of the surveys and t is the time since the first year. The estimated rate of population decline ($\hat{\delta}$) was lowest in the northern regions and highest in the southern regions. If only data since 1960 are considered, the population decline on St. Pierre Bank, Sydney Bight, and Banquereau Bank is similar to that in the southernmost regions (that is, Gulf of Maine, Georges Bank, and southern New England). The standard error (SE) of $\hat{\delta}$ is provided.

References and Notes

1. M. J. Holden, *Rapp. P.-V. Reun. Cons. Int. Explor. Mer* **164**, 360 (1973).
2. _____, in *Sea Fisheries Research*, F. R. H. Jones, Ed. (Elek Science, London, 1974), pp. 117-137.
3. P. A. Walker and J. R. G. Hislop, *ICES J. Mar. Sci.* **55**, 392 (1998).
4. E. D. Anderson, in *Elasmobranchs as Living Resources: Advances in the Biology, Ecology and the Status of the Fisheries*, H. L. Pratt Jr., S. H. Gruber, T. Taniuchi, Ed. (U. S. Department of Commerce, Washington, DC, 1990), pp. 473-484.
5. S. A. Murawski and J. S. Doine, *J. Northwest Atl. Fish. Sci.* **14**, 79 (1992).
6. M. J. Fogarty and S. A. Murawski, *Ecol. Appl.* **8** (suppl. 1), S6 (1998).
7. H. B. Bigelow and W. C. Schroeder, *Fishes of the Western North Atlantic, Part 2* (Sears Foundation for Marine Research, New Haven, CT, 1953).
8. In May 1929 in the southern New England region, barndoor skates were captured at an average rate of seven per tow (7). On Georges Bank in 1951, the average capture rate for barndoor skates was as high as 21 per tow [H. B. Bigelow and W. C. Schroeder, *Fish. Bull.* **53**, 61 (1953)].
9. G. R. Huntsman, *Fisheries* **19**, 8 (1994).
10. K. Brander, *Nature* **290**, 48 (1981).
11. Research vessel survey data on the barndoor skate have been assembled from the southern Grand Bank, at the northern limit of the range, to southern New England. From 1971 in Canadian waters and 1963 in

the United States, annual research surveys have been conducted with a stratified random survey design [W. G. Doubleday and D. Rivard, Eds., *Bottom Trawl Surveys* (Department of Fisheries and Oceans, Ottawa, Canada, 1981)]. These surveys cover a range of depths from 50 to 400 m and are designed to provide unbiased estimates of abundance. Before 1971, a fixed location survey along line transects was used off the coasts of Newfoundland and Nova Scotia. These surveys were converted to the stratification scheme with the latitude, longitude, and depth of the tow. The earlier surveys were conducted primarily during the day, and, because skates are caught in substantially higher proportions at night, day catches would underestimate the true abundance (J. M. Casey and R. A. Myers, *Can. J. Fish. Aquat. Sci.*, in press). To account for this difference in diel catchability, we converted half of the catches before 1970 to night catches using a factor of 2.08. These surveys would then be comparable with those after 1970 in which day and night tows were conducted in roughly equal proportion. Biomass estimates were based on the stratified random design [W. J. Cochran, *Sampling Techniques* (Wiley, New York, 1977)]. Estimates of absolute abundance were obtained by dividing the biomass estimates by a factor of 0.15, if 10% of the skates in the path of the trawl are caught during the day [R. L. Edwards, in *The Future of the Fishing Industry of the United States*, D. Gilbert, Ed. (University of Washington, Seattle, WA, 1968), pp. 52–60] and 20% of the population would be caught at night. Neither of these conversions affects the observed biomass trends.

12. Because barndoor skates have only been caught in research surveys of Browns Bank and Georges Bank in the past 20 years, we examined the statistical power of detecting barndoor skate given a specific low abundance level. On St. Pierre Bank, 504 of 1075 research survey tows were conducted over the past 25 years at depths where, historically, barndoor skates were commonly found (200 to 400 m). We assume that the number n of barndoor skates caught in one tow has a negative binomial distribution, with a probability of $[\Gamma(k+n)/\Gamma(k)n!][\mu^k k! / (\mu+k)^{k+n}]$, where μ is the mean catch per tow and the constant k is the inverse of aggregation. Then, with a theoretical low mean abundance of 500 individuals in a 7368-km² area (the tow sample area is 0.05 km², and the probability of catching a barndoor skate if encountered is 0.15) and $k = 0.5$, the probability of not detecting a barndoor skate in 504 tows is 0.77. Greater aggregation (that is, lower k) results in only small changes in this number. If 1000 individuals remained, the probability of not detecting a barndoor skate decreases to 0.6.
13. W. Templeman, *Marine Resource of Newfoundland* (Bulletin Fisheries Research Board of Canada, Ottawa, Canada, 1966).
14. ———, *ICNAF Res. Bull.* **13**, 31 (1978).
15. R. A. Myers, G. Mertz, S. Fowlow, *Fish. Bull.* **95**, 762 (1997).
16. The relation between the weight of the young at hatching, w , and the maximum estimate of fecundity is $10.9w^{-0.45}$ (1).
17. For the barndoor skate population to be self-sustaining, the survival to age x , l_x , must be sufficient such that $\sum l_x m_x = 1$, where m_x is the number of eggs developing into females from individuals at age x . If survival, P , is constant such that $l_x = P^x$, then $P^a m_0 \sum_{x=0}^{\infty} P^x = 1$, where the survival to the age of maturity a is P^a and the fecundity in each year is m_0 . Summing the geometric series, $P^a m_0 [1/(1-P)] = 1$. The instantaneous mortality rate is $\log(P)$. If the mortality of juveniles is twice that of adults, the instantaneous mortality required to drive the barndoor skate to extinction is 0.45.
18. This age estimate is based on the relation between somatic growth rate and temperature and a comparison of age at maturity of cod that occur at the same locations (15).
19. R. A. Myers, J. A. Hutchings, N. J. Barrowman, *Mar. Ecol. Prog. Ser.* **138**, 293 (1996).
20. N. Shackell and J. Lien, in *Marine Protected Areas and Sustainable Fisheries*, N. L. Shackell and J. H. Martin

Willison, Eds. (Science and Management of Protected Areas Association, Nova Scotia, Canada, 1995), pp. 21–30.

21. We thank the Department of Fisheries and Oceans at St. John's and Halifax, Canada, and the National Marine Fisheries Service at Woods Hole, MA, for providing the data used in this analysis and B. Atkin-

son, K. Frank, J. E. Simon, M. Fogarty, S. Murawski, and J. Hutchings for helpful discussions. This research was initiated while R.A.M. was a visiting fellow at the Centre for Population Biology, Imperial College at Silwood Park.

17 February 1998; accepted 18 June 1998