

By-catch of Harbour Porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy Gillnet Fishery, 1998–2001

E.A. Trippel and T.D. Shepherd

Fisheries and Oceans Canada
Biological Station
531 Brandy Cove Road
St. Andrews, NB
E5B 2L9

2004

**Canadian Technical Report of
Fisheries and Aquatic Sciences 2521**



Fisheries and Oceans Canada Pêches et Océans
Canada Canada

Canada

Canadian Technical Report of
Fisheries and Aquatic Sciences 2521

2004

**BY-CATCH OF HARBOUR PORPOISE (*PHOCOENA PHOCOENA*) IN
THE LOWER BAY OF FUNDY GILLNET FISHERY, 1998–2001**

by

E.A. Trippel and T.D. Shepherd

Fisheries and Oceans Canada
Biological Station, 531 Brandy Cove Road,
St. Andrews, New Brunswick, Canada, E5B 2L9

This is the two hundred and fifty-second Technical Report
of the Biological Station, St. Andrews, NB

© Her Majesty the Queen in Right of Canada, 2004.
Cat. No.Fs97-6/2521E ISSN 0706-6457

Correct citation for this publication:

Trippel, E.A., and T.D. Shepherd. 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery, 1998-2001. Can. Tech. Rep. Fish. Aquat. Sci. 2521: iv + 33 p.

ABSTRACT

Trippel, E.A., and T.D. Shepherd. 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery, 1998-2001. Can. Tech. Rep. Fish. Aquat. Sci. 2521: iv +33 p.

The most serious threat to the status of harbour porpoise (*Phocoena phocoena*) is incidental mortalities caused by entanglement in fishing gear. As part of an ongoing study to evaluate the utility of mitigation techniques to reduce such harbour porpoise mortalities, observers were placed on-board vessels participating in the lower Bay of Fundy demersal gillnet fishery from 1998-2001. The number of vessels participating in the fishery declined from 22 in 1998 to 13 in 2001. Despite this decline, total annual effort remained relatively stable. Over the 4-yr period, a total of 52 porpoise mortalities were observed, all but four of which were in the Swallowtail region off Grand Manan, New Brunswick. For the Bay of Fundy, total estimated by-catch was 38, 32, 28, and 73 porpoises from 1998-2001, respectively. A lack of spatial and temporal observer coverage did not allow us to generate estimates of variance in by-catch estimates. In all years, the majority of the estimated porpoise by-catch occurred in the Swallowtail region. In general, by-catch was highest in July except in 1999 when it was highest in August. The increase in by-catch seen in 2001 was due to increased catch rates rather than increased effort. While the Canadian by-catch in 2001 appears low (73 porpoises), it was nearly as high as the U.S. by-catch (80 porpoises) which has a much larger fishery. Given that the Canadian by-catch appears to represent a significant source of mortality, observer coverage should be better allocated both spatially and temporally in the future so that more reliable by-catch estimates and their associated variance can be estimated.

RÉSUMÉ

Trippel, E.A., and T.D. Shepherd. 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery, 1998-2001. Can. Tech. Rep. Fish. Aquat. Sci. 2521: iv +33 p.

La plus sérieuse menace au statut des marsouins communs (*Phocoena phocoena*) concerne les mortalités accidentelles causées par l'empêchement dans les engins de pêche. Dans le cadre d'une étude visant l'évaluation de l'utilité de techniques mitigatives pour réduire ce type de mortalités chez le marsouin commun, des observateurs ont été embarqués à bord de bateaux participant à la pêche au poisson de fond à l'aide de filets maillants dans la Baie de Fundy entre 1998–2001. Le nombre de bateaux participant à cette pêche a diminué de 22 en 1998 à 13 en 2001. Malgré ce déclin, l'effort annuel total est demeuré relativement stable. Au cours de ces 4 années, un total de 52 mortalités de marsouins ont été observées, toutes sauf 4 étant survenues dans la région Swallowtail au large de Grand Manan, Nouveau-Brunswick. Pour la Baie de Fundy, les prises accidentelles totales de 1998 à 2001 ont été estimées à 38, 32, 28, et 73 marsouins, respectivement. Un manque de couverture spatiale et temporelle des observateurs en mer n'a pas permis d'estimer la variance autour de ces estimations de prises accidentelles. Pour chacune des années, la

majorité des prises accidentelles de marsouins sont survenues dans la région Swallowtail. En général, les prises accidentelles ont été les plus élevées en juillet, sauf en 1999 où elles ont été les plus élevées en août. L'augmentation des prises accidentelles observée en 2001 était liée à l'augmentation du taux de capture plutôt qu'à une augmentation de l'effort. Bien que les prises accidentelles canadiennes puissent sembler faibles en 2001 (73 marsouins), elles étaient presque aussi élevées que les prises accidentelles des États-Unis (80 marsouins) qui mène une beaucoup plus importante pêcherie. Considérant que les prises accidentelles canadiennes semblent constituer une source significative de mortalité, la couverture par les observateurs en mer devrait être mieux distribuée dans le temps et l'espace dans le futur de manière à obtenir des estimations fiables des prises accidentelles et de la variance de ces estimations.

INTRODUCTION

Incidental mortalities of harbour porpoise (*Phocoena phocoena*) occur in demersal gillnets throughout most of its range (Gaskin 1984; Read and Gaskin 1988; Fontaine et al. 1994; Perrin et al. 1994; Trippel et al. 1996a; Murray et al. 2002). Mortalities have occurred in the Bay of Fundy since the regular use of gillnets began in the 1960's (Gaskin 1992). Beginning in 1993, observer programs have been conducted each summer in the Bay of Fundy to collect data related to by-catch of harbour porpoise. Observed mortalities were combined with data on fishing effort to yield annual estimates of by-catch that ranged from 20 to 424 animals for the period 1993-97 (Trippel et al. 1996a, 1999; Table 1).

Harbour porpoise occurring in the Bay of Fundy (Fig. 1) form part of a population of ~90,000 animals (Waring et al. 2001) that undergo seasonal north-south migrations along the U.S.-Canada coast from North Carolina to Nova Scotia (Read and Westgate 1997). Analysis of mitochondrial DNA indicates this is a discrete population (Wang et al. 1996; Rosel et al. 1999) and is commonly referred to as the Bay of Fundy/Gulf of Maine population (BOF/GOM). The U.S. National Marine Fisheries Service (NMFS) has provided estimates of porpoise by-catch for the Gulf of Maine and waters off the Mid-Atlantic States that since 1990 ranged from 80 to 2,900 animals (Table 1). Two other harbour porpoise populations exist in eastern Canada and are referred to as the Gulf of St. Lawrence and Newfoundland-Labrador populations (Rosel et al. 1999). The BOF/GOM population is considered by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) to be of "special concern" (COSEWIC 2003). In January 1993, NMFS proposed listing the BOF/GOM harbour porpoise as Threatened under the U.S. *Endangered Species Act* (NMFS 1993). In January 1999, after 6 yr of action NMFS supported removal of this population from the candidate list citing bilateral by-catch reduction programs and recent population estimates as evidence for non-threatened status (NMFS 1999, 2001). The harbour porpoise is classified as vulnerable in the IUCN Red List (IUCN 2002).

By-catch mitigation research has been underway in the Bay of Fundy since 1994 (Lien et al. 1995; Trippel et al. 1996a, 2003). Acoustic pingers (Dukane Netmark 1000) were field tested and shown to be effective in reducing by-catch by 68% in 1996 and 85% in 1997 (Trippel et al. 1999). A new type of gillnet developed to be more acoustically reflective than 100% nylon netting has been tested since 1998 in the Bay of Fundy. Testing of this nylon barium-sulphate gillnet in 1998 and 2000 led to zero by-catch in reflective nets and 12 porpoises in control nets (Trippel et al. 2003). Field-testing of a similar version containing iron oxide particles also led to a significant reduction in porpoise by-catch in the North Sea (Larsen et al. 2002).

In Canada, the DFO Maritimes Region has developed a Harbour Porpoise Conservation Strategy and has set a maximum take of 110 harbour porpoises per year for the Bay of Fundy, a mortality believed to have a negligible effect on the growth potential of this transboundary population (DFO 1995). In cooperation with the U.S., Canada annually provides preliminary estimates of by-catch for the Bay of Fundy to complete U.S. estimates of the Potential Biological Removal Rate (PBR), which is a conservative

estimate of the allowable removal level of animals from the population (Wade 1998). Most recently, the PBR level has been set at 747 porpoises per year for the entire population (Bay of Fundy, Gulf of Maine and Mid-Atlantic States; Waring et al. 2001). Canadian calculations are conducted as the season progresses to alert science and resource managers to by-catches approaching 110 animals, so that a closure of sensitive areas to gillnet fishing can be implemented (DFO 1995). Although preliminary within-season values of by-catch are made available, it is necessary to periodically produce peer-reviewed estimates of by-catch. The by-catch estimation procedures are at times complicated as they must account for differential use of control and mitigative gear and corresponding fishing effort. Moreover, observer coverage has been absent in some areas known to experience by-catch, thus necessitating additional assumptions that make it difficult to establish error limits about estimates. However, in several key fishing areas that are receiving observer coverage, it is common to have a very high percentage of the fleet covered by observers (>50%, Trippel et al. 1996a). The most recent peer-reviewed estimate of porpoise by-catch for the Bay of Fundy gillnet fishery was for the 1997 season, although a preliminary estimate was given for 1998 (DFO 1998). In this report, we have updated the porpoise by-catch analysis for the Bay of Fundy for the period 1998-2001 by addressing the following objectives.

- 1) Determine the number of fishing vessels and effort on different fishing grounds both seasonally and annually.
- 2) Report the number of observed porpoise mortalities in gillnets and plot their spatial distribution.
- 3) Estimate seasonal, spatial and total annual by-catch.

METHODS

DATA SOURCE

From 1998 until 2001, DFO and others were part of an effort to develop and test the effectiveness of a barium-sulphate filled gillnet mesh in reducing harbour porpoise by-catch (Trippel et al. 2003). As a part of this study, observers were placed on board gillnet fishery vessels in the lower Bay of Fundy in order to quantify porpoise and groundfish catches in both regular nylon mesh nets and barium-sulphate mesh nets. Observer coverage in 1998 was augmented by fishing vessels that operated without an observer but who participated in a voluntary reporting program on the effectiveness of the two types of gillnet mesh. Field coordination of the observer program was provided by the Grand Manan Fishermen's Association (1998, 2000, and 2001) and Javitech Ltd. (1999).

OBSERVED PORPOISE MORTALITIES

There is evidence that barium-sulphate gillnet mesh lowers by-catch of harbour porpoise (Trippel et al. 2003). Observed vessels during the 1998-2001 period most often used both nylon-mesh strings and barium-sulphate mesh strings on any one trip, while unobserved vessels exclusively used nylon-mesh strings. Because of this, observed data were

summarized at the string level to avoid the potentially confounding effect of mesh type. Data were first partitioned by year, fishing area (Fig. 2) and season (2-wk periods). A small number of observations did not include location data (4%). These observations were assigned to a fishing location based on known locations fished by vessels from the same port of origin during that year and season. Catch rates of harbour porpoise (mean catch per string) were calculated for each year, fishing location, and season combination. A typical fishing trip by a lower Bay of Fundy gillnet vessel involves fishing five strings. A string is defined as three approximately 100-m long webs that are fished as a unit for approximately 24 h before retrieval. To facilitate the application of observed catch rates to unobserved data, catch rates per string were multiplied by five in order to derive an estimate of porpoise by-catch per trip.

ESTIMATING CATCH RATES

Spatial and temporal observer coverage in the dataset was generally low except at Swallowtail (Table 3). Since 1994, observed coverage has been in place at Swallowtail for each year except 1999. Because of this, Swallowtail catch rates were used as a “standard” against which unknown catch rates for other locations could be estimated. While the fishery begins in June, effort is typically very low and there has never been any porpoise by-catch reported in June in the lower Bay of Fundy. Because of this, it was assumed that porpoise catch rates in June of each year were zero. Observer coverage extended past September 15 only in 1999 at the Wolves during which no porpoises were captured. Because of this, it was assumed that porpoise catch rates after September 15 of each year were zero. In a number of fishing grounds (i.e., Grand Manan Banks, Head & Horns, and Digby Neck; Fig. 1) there have not been any instances of observed by-catch in the gillnet fishery. The by-catches in these areas were assumed to be zero in all years.

In order to estimate catch rates when no observer coverage was present, ratios were calculated between the catch rates of each fishing location and Swallowtail using data from the years 1994–2001, i.e., where paired observations during a season existed between Swallowtail and a particular fishing area, the ratio of its catch rate to Swallowtail was calculated. The mean of these ratios from all pairs for a particular fishing area represented the mean ratio of its catch rate relative to Swallowtail. When observer coverage was absent for a particular fishing location/season/year combination, it was estimated by prorating the associated Swallowtail catch rate by the mean ratio relative to Swallowtail.

In 1999, observer coverage was absent from Swallowtail but was extensive at the Wolves. In order to estimate Swallowtail catch rates, seasonal catch rates from the Wolves were multiplied by the reciprocal of the Wolves mean ratio to Swallowtail from other years. In some years, seasonal catch rates were missing from Swallowtail. They were estimated by using the seasonal mean ratio of catch rates, using August 1-15 as a standard in a manner analogous to fishing location mean ratios. August -15 was selected as a standard since observer coverage was most complete for this period.

ESTIMATING PORPOISE BY-CATCH

Observed effort (trip level) was first subtracted from reported effort (trip level) in order to derive unobserved effort. Reported effort was extracted from the DFO commercial fishery database which includes information on location fished and landings for each trip. Catch rates for each year/location/season were multiplied by the unobserved effort of the associated year/location/season combination in order to estimate unobserved porpoise by-catch. Total estimated by-catch was obtained by adding the observed by-catch to the estimated unobserved by-catch.

RESULTS

THE FISHERY

From 1998–2001, the number of vessels participating in the fishery decreased from 22 in 1998 to 13 in 2001 (Table 2). Of the vessels participating, most fished only through July and August. Fishing effort by at least some participating vessels increased over the study period since the total number of trips reported by participating vessels fell only modestly from 276 trips in 1998 to 257 trips in 2001 (Table 3). The spatial distribution of fishing effort appeared to shift between 1998 and 2001. In all 4 yr the majority of the effort was concentrated on Swallowtail (Table 3). In 1998, Gravelly Bulkhead was secondary to Swallowtail in effort reported. In subsequent years, effort at Gravelly Bulkhead decreased while it increased at the Wolves (Table 3). By 2001, the Wolves was secondary to Swallowtail in fishing effort.

OBSERVED PORPOISE MORTALITIES

Over the entire 4-yr period, 52 porpoise captures were observed (Table 4). Observed porpoise by-catch was low from 1998-2000 and relatively high in 2001 (Table 5). From 1998-2001 porpoise by-catch was observed in three areas: Swallowtail, the Wolves and Gravelly Bulkhead (though minimal observer coverage occurred elsewhere). All but four porpoise captured were observed at Swallowtail (Table 5; Fig. 2). Two of the porpoise catches at the Wolves were in mixed strings (i.e., strings with both 100% nylon-mesh webs and barium-sulphate mesh webs). In both cases, porpoises were captured in 100% nylon-mesh webs. General trends in observed catch rates per string (Table 6) were similar to observed catch rates per trip (Table 5). Not unexpectedly, when catch rates per string were adjusted to the trip level using five strings per trip (Table 7) they became much higher than the observed catch rates per trip (Table 5).

DEVELOPMENT OF PRO-RATED CATCH RATES TO ESTIMATE BY-CATCH

The observer database encompasses the years 1993-2001. Data from 1993 were not used in this analysis since information on the geographic location of fishing effort was absent. Rather, 1993 included data on port of origin (Trippel et al. 1996a). During the period from 1994-2001, porpoise catches were observed at Swallowtail, Wolves, Gravelly Bulkhead, Head Harbour and the Channel (Table 8; Fig. 2). At no time during this period was there total spatial and temporal observer coverage. Relative to Swallowtail, historical catch rates

in Head Harbour and Gravelly Bulkhead were high while those at the Wolves and the Channel were low (Table 9). A considerable amount of uncertainty existed for some estimates of mean ratio catch rates (e.g., Gravelly Bulkhead) due to the low number of seasonal pairs available. Seasonal catch rates at Swallowtail were highest from July 1-31 (Table 9). Using Swallowtail as a standard allowed us to fill in unknown catch rates for all year/location/season combinations (Table 10).

PORPOISE BY-CATCH

Total estimated porpoise by-catch (observed by-catch plus estimated unobserved by-catch) for the lower Bay of Fundy was 38, 32, 28, and 73 porpoises for 1998-2001, respectively (Fig. 3). The 2001 by-catch (73 porpoises) was lower than by-catch reported prior to 1996, but higher than those reported between 1996 and 2000 (Fig. 3). Total estimated by-catch was highest at Swallowtail during all years and accounted for between 43.8% and 75.3% of total estimated by-catch annually (Table 11; Fig. 4). Effort in the Channel was not sufficient to generate estimated unobserved by-catch. Total estimated by-catch in the other areas was generally low relative to Swallowtail. By-catch was generally highest in either early or late July except 1999 when almost all of the total estimated by-catch occurred in late August (Table 11) and observer effort was dedicated exclusively to the Wolves (Table 5). The increased by-catch in 2001 appeared to be due to increased catch rates (Fig. 5). Total catch rates calculated from observed by-catch and total estimated by-catch showed a similar trend where it was relatively low from 1998-2000 and subsequently increased considerably in 2001 (Fig. 5).

DISCUSSION

The annual by-catch estimates for the period 1998-2001 have remained below the 110 animals set as a cap by the DFO Maritimes Region's Harbour Porpoise Conservation Strategy (DFO 1995). The most recent BOF/GOM population estimate was reported to be 89,700 animals (survey conducted in 1999; Waring et al. 2001), and consequently, this level of mortality is not considered to be high relative to population size (0.12%). Reduced quotas and shortened fishing seasons under restrictive groundfish management plans are primarily responsible for the maintenance of annual by-catches in the Bay of Fundy of less than 110 porpoises since the mid-1990s. The cumulative mortality in both Canada and the U.S. for this population is shown in Table 1. The by-catch for the Bay of Fundy in 2001 was surprisingly nearly equivalent to the entire U.S. by-catch for this population. This is partly due to the high by-catch rate observed in the Swallowtail area, Bay of Fundy, from July 16-31 of 2001: 0.625 porpoises/trip in 100% nylon nets and 0.360 porpoises/trip in barium-sulphate nets. Only in 1994 was a similar by-catch rate observed in control gear with a mean of 0.61 porpoises/trip occurring from August 15-31 in the Swallowtail area (Trippel et al. 1996a). By-catch rate was also high in 1993 by vessels fishing out of Grand Manan (3.20 porpoises/trip; Trippel et al. 1996a). The high abundance of porpoises in the BOF/GOM population, the annual variability in the portion of the population that enters the Bay of Fundy, and the resulting potentially high by-catch rates indicates the importance of annual monitoring of by-catch in the Bay of Fundy.

This report highlights the need for improved observer coverage. Fishing in areas prone to high porpoise by-catch rates (Wolves and Head Harbour fishing grounds) by vessels departing from Campobello Island did not receive annual observer coverage. This necessitated pro-rating of known by-catch rates to areas of unknown by-catch rates (e.g., rate in one area such as Swallowtail to another area), presumably leading to inaccuracies and uncertainty in the number of estimated mortalities. Moreover, only rarely over the previous 10 yr (e.g., 1994) did widespread observer coverage exist for the more distant areas of the Bay of Fundy accessed by both New Brunswick and Nova Scotia gillnetters (e.g., Grand Manan Basin, Northeast Bank). Nova Scotia gillnetters in the Bay of Fundy were rarely covered and were assumed not to catch porpoises (based on zero by-catch during sporadic observer coverage). However, porpoise sightings are frequent in these areas during population surveys (Palka 1995a) suggesting entanglements in Nova Scotia fishing gear may be occurring. It is recommended that another year of widespread observer coverage be conducted in the Bay of Fundy and south-west Nova Scotia in order to establish whether zero by-catch is a reasonable assumption for these areas and to re-evaluate the level of pro-rating currently applied among areas.

The by-catch patterns in September 2001 alert us to a number of important issues. Not since 1994 have estimates been made of bi-weekly by-catch through the entire fishing season (i.e., early July–mid September). In 2001, by-catch rates were lower in August than in either July or September. In contrast, 1994 by-catch rates were highest in August. Annual changes in seasonal variation of by-catch of porpoise in the Bay of Fundy seem to occur and consequently closing the fishery every year during a fixed 2-wk period may not necessarily reduce by-catch significantly. Earlier analyses led to a possible recommendation to reduce by-catch by simply closing the fishery in sensitive areas from August 16-30 (in 1994 total by-catch would have been reduced by 38% by this 2-wk closure; Trippel et al. 1996a). However, this 2-wk closure of the Swallowtail and Wolves areas would not have had a significant impact in 2001 (5.5% of total estimated by-catch). It should also be noted that the relatively high by-catch rate in September 2001 should reduce confidence in our assumption that no by-catch occurred after September 15 of each year.

Alternatively, enforced use of mitigative gear could be a more suitable management option to explore. To date, it is not clear from experimental tests which is the most suitable form of mitigation (acoustic pingers vs. reflective nets). Results on the efficacy of reflective gillnets in 2001 (nylon: 0.16 porpoises/trip, barium-sulphate: 0.11 porpoises/trip) were not as promising as earlier research in 1998 (nylon: 0.08 porpoises/trip, barium-sulphate: 0 porpoises/trip) and 2000 (nylon: 0.05 porpoises/trip, barium-sulphate: 0 porpoises/trip). In 1999 mixed webs were used flawing the experimental design. The change in effectiveness may be associated with an unusually high abundance of harbour porpoises in the Bay of Fundy in the summer of 2001 (entrapment of porpoises in weirs was among the highest recorded – 1998 = 34, 1999 = 93, 2000 = 20, 2001 = 312, 2002 = 53, and 2003 = 31; source: Grand Manan Whale & Seabird Station). By-catch rates in gillnets (100% nylon) in 2001 were the second highest in the period from 1994-2001 (Table 8; July 16-31, 2001

at Swallowtail: 0.625 porpoises/trip, Aug 16-31 1994 at Head Harbour and July 1-15, 1998 at Gravelly Bulkhead: 1.0 porpoises/trip).

Abundance of porpoises in the Bay of Fundy is positively correlated with abundance of Atlantic herring (*Clupea harengus*) (Palka 1995b; Trippel et al. 1999), their principal prey (Recchia and Read 1989). We speculate the aversive behaviour of porpoises around gillnets in 2001 may have been altered and possibly diminished from earlier test years (Trippel et al. 2003) as porpoise could have been exhibiting aggressive swimming and diving behaviour while foraging resulting in reduced alertness to barriers such as gillnets. Because of this, integration of herring abundance data, from fishery sources, may assist in the prediction of regionally high porpoise by-catch levels. On the other hand, the effectiveness of reflective netting may be inherently lower than previously reported (Trippel et al. 2000; Larsen et al. 2002; Trippel et al. 2003). Consequently, it is recommended that further field-testing of barium-sulphate gillnets be undertaken in the Bay of Fundy under various levels of regional porpoise abundance. The present period of reduced groundfish quotas provides a good opportunity to conduct these field trials without incurring large losses of animals.

ACKNOWLEDGEMENTS

We thank the Grand Manan Fishermen's Association and Javitech Ltd. for providing field project coordination and data collection aboard the fishing vessels. Don King and Norm Holy of Atlantic Gillnet Supply collaborated on provision of the barium-sulphate gillnets for field testing. By-catch coverage and experimental testing was supported by Fisheries and Oceans Canada and the U.S. National Marine Fisheries Service. We also thank Jack Lawson and Véronique Lesage for their constructive comments on an earlier version of the manuscript. Michele Saunders and Brenda Best assisted with the final stages of manuscript preparation.

REFERENCES

- Committee on the Status of Endangered Wildlife in Canada (COSEWIC) 2003. Canadian Species at Risk. (Available from COSEWIC Secretariat, Ottawa, ON).
- Department of Fisheries and Oceans (DFO) 1995. Harbour porpoise conservation strategy for the Bay of Fundy. (Available from Department of Fisheries and Oceans, Resource Management Branch, P.O. Box 550, Halifax, NS B3J 2S7, Canada).
- Department of Fisheries and Oceans (DFO) 1998. Harbour porpoise by-catch in the lower Bay of Fundy gillnet fishery. DFO Maritimes Regional Status Report 98/7E. (Available from Maritimes Regional Advisory Process, Department of Fisheries and Oceans, P.O. Box 1006, Stn. B203, Dartmouth, NS B2Y 4A2, Canada).

- Fontaine, P.-M., Barrette, C., Hammill, M.O., and Kingsley, M.C.S. 1994. Incidental catches of harbour porpoises (*Phocoena phocoena*) in the Gulf of St. Lawrence, and the St. Lawrence River estuary, Quebec, Canada. *Int. Whaling Comm. Rep. Comm. (Spec. Issue) 15*: 159-163.
- Gaskin, D.E. 1984. The harbour porpoise *Phocoena phocoena* (L.): regional populations, status, and information on direct and indirect catches. *Int. Whaling Comm. Rep. Comm. 34*: 569-586.
- Gaskin, D.E. 1992. Status of the harbour porpoise, *Phocoena phocoena*, in Canada. *Can. Field-Nat. 196*: 36-54.
- IUCN 2002. 2002, IUCN redlist of threatened species. www.redlist.org Downloaded January 27, 2003.
- Larsen, F., Eigaard, O.R., and Tougaard, J. 2002. Reduction of harbour porpoise by-catch in the North Sea by high-density gillnets. *Int. Whaling Comm. SC/54/SM30*. 12 p.
- Lien, J., Hood, C., Pittman, D., Ruel, P., Borggard, D., Chisholm, C., Weisner, L., Mahon, T., and Mitchell, D. 1995. Field tests of acoustic devices on groundfish gillnets: assessment of effectiveness in reducing porpoise by-catch. *In* *Sensory systems of aquatic mammals*. Edited by R.A. Kastelein, J.A. Thomas and P.E. Nachtigall. De Spil Publishers, Woerden, The Netherlands. pp. 349-364.
- Marine Mammal Commission. 2002. Marine Mammal Commission Annual Report to Congress – 2001. Marine Mammal Commission, Bethesda, Maryland.
- Murray, K.T., Read, A.J., and Solow, A.R. 2002. The use of time/area closures to reduce bycatches of harbour porpoises: lessons from the Gulf of Maine sink gillnet fishery. *J. Cetacean Res. Manag. 2*: 135-141.
- National Marine Fisheries Service. 1993. Proposed listing of Gulf of Maine population of harbor porpoise as threatened under the Endangered Species Act. *Fed. Reg. 58*: 3108-3120. January 07, 1993.
- National Marine Fisheries Service. 1999. Listing of Gulf of Maine/Bay of Fundy population of harbor porpoise as threatened under the Endangered Species Act. *Fed. Reg. 64*: 465-471. January 05, 1999.
- National Marine Fisheries Service. 2001. Status review of the Gulf of Maine/Bay of Fundy population of harbor porpoise under the Endangered Species Act. *Fed. Reg. 64*: 40176-40187. August 02, 2001.
- Palka, D. 1995a. Abundance estimate of the Gulf of Maine harbor porpoise. *Int. Whaling Comm. Rep. Comm. (Spec. Issue) 16*: 27-50.

- Palka, D. 1995b. Influences on spatial patterns of Gulf of Maine harbor porpoises. *In* Whales, seals, fish and man. Edited by A.S. Blix, L. Walløe and Ø. Ulltang. Elsevier Science B.V., The Netherlands. pp. 69-75.
- Palka, D. 1997. Gulf of Maine Harbor Porpoise By-catch. Prepared for the Gulf of Maine Harbor Porpoise Take Reduction Team Meeting, December 16-17, 1997. National Marine Fisheries Service, Woods Hole, Massachusetts.
- Perrin, W.F., Donovan, G.P., and Barlow, J. 1994. Report of the workshop on mortality of cetaceans in passive fishing nets and traps. *Int. Whaling Comm. Rep. Comm.* 15: 6-57.
- Read, A.J., and Gaskin, D.E. 1988. Incidental catch of harbour porpoises by gill nets. *J. Wildl. Manag.* 52: 517-523.
- Read, A.J., and Westgate, A.J. 1997. Monitoring the movements of harbour porpoises (*Phocoena phocoena*) with satellite telemetry. *Mar. Biol.* 130: 315-322.
- Recchia, C.R., and Read, A.J. 1989. Stomach contents of harbour porpoises, *Phocoena phocoena* (L.), from the Bay of Fundy. *Can. J. Zool.* 67: 2140-2146.
- Rosel, P.E., France, S.C., Wang, J.Y., and Kocher, T.D. 1999. Genetic structure of harbour porpoise *Phocoena phocoena* populations in the northwest Atlantic based on mitochondrial and nuclear markers. *Mol. Ecol.* 8: S41-S54.
- Trippel, E.A., Holy, N.L., Palka, D.L., Shepherd, T.D., Melvin, G.D., and Terhune, J.M. 2003. Nylon barium sulphate gillnet reduces porpoise and seabird mortality. *Mar. Mamm. Sci.* 19: 240-243.
- Trippel, E.A., Palka, D., Potter, D., Merrick, R., Holy, N.L., King, D., Conway, J.D., and Shepherd, T. 2000. Testing of acoustically reflective gillnets as a management tool to reduce incidental mortalities of harbour porpoise in the Northwest Atlantic gillnet fisheries. *Int. Whaling Comm. SC/52/SM26.* 9 p.
- Trippel, E.A., Strong, M.B., Terhune, J.M., and Conway, J.D. 1999. Mitigation of harbour porpoise (*Phocoena phocoena*) by-catch in the gillnet fishery in the lower Bay of Fundy. *Can. J. Fish. Aquat. Sci.* 56: 113-123.
- Trippel, E.A., Wang, J.Y., Strong, M.B., Carter, L.S., and Conway, J.D. 1996a. Incidental mortality of harbour porpoise (*Phocoena phocoena*) by the gill-net fishery in the lower Bay of Fundy. *Can. J. Fish. Aquat. Sci.* 53: 1294-1300.
- Trippel, E.A., Strong, M.B., Hood, C., Richter, C., and Lien, J. 1996b. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery in 1995. *DFO Atl. Fish. Res. Doc.* 96/110. 13 p. (Available from the Canadian Stock Assessment Secretariat, Ottawa, ON).

- Wade, P.R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Mar. Mamm. Sci.* 14: 1-37.
- Wang, J.Y., Gaskin, D.E., and White, B.N. 1996. Mitochondrial DNA analysis of harbour porpoise, *Phocoena phocoena*, subpopulations in North American waters. *Can. J. Fish. Aquat. Sci.* 53: 1632-1645.
- Waring, G.T., Quintal, J.M., and Swartz, S.L. (ed.). 2001. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments – 2001. NOAA Technical Memorandum NMFS-NE-168. Northeast Fisheries Science Center, Woods Hole, MA.
- Westgate, A.J., Read, A.J., Berggren, P., Koopman, H.N., and Gaskin, D.E. 1995. Diving behaviour of harbour porpoises, *Phocoena phocoena*. *Can. J. Fish. Aquat. Sci.* 52: 1064-1073.

Table 1. Estimates of harbour porpoise by-catch in demersal gillnet fisheries in the Bay of Fundy, New England, and off the U.S. mid-Atlantic states from 1990-2001. Numbers in parentheses for U.S. data represent 95% confidence intervals of by-catch estimates. Numbers in parentheses for Bay of Fundy data represent ± 1 standard error and 95% confidence limits for 1993 and 1994, respectively.

Year	New England ¹	Bay of Fundy ²	U.S. Mid-Atlantic ³	Other ⁴	Total
1990	2900 (1500 – 5000)	-	-	-	2900
1991	2000 (1000 – 3800)	-	-	-	2000
1992	1200 (800 – 1700)	-	-	-	1200
1993	1400 (1000 – 2000)	424 (200 – 648)	-	-	1824
1994	2100 (1400 – 2900)	101 (80 – 122)	-	-	2201
1995	1400 (900 – 2500)	87	103 (11 – 254)	-	1590
1996	1200 (800 – 1800)	20	311 (162 – 567)	-	1531
1997	782 (501 – 1208)	43	572 (296 – 1071)	-	1397
1998	332 (170 – 728)	38	446 (294 – 894)	-	816
1999	270 (78 – 364)	32	53 (3 – 98)	19	374
2000	570 (169 – 924)	28	21 (1 – 53)	1	620
2001	51 (2 – 166)	73	26 (1 – 83)	3	153

¹ 1990-1996 from Palka 1997, 1997-1999 from Waring et al. 2001, 2000 from Marine Mammal Commission 2002, 2001 from D.Laist, Marine Mammal Commission.

² 1993-1994 from Trippel et al. 1996a, 1995 from Trippel et al. 1996b, 1996-1997 from DFO 1998, 1998-2001 from this study.

³ 1995-1996 from Palka 1997, 1997-2000 from Marine Mammal Commission 2002, 2001 from D. Laist, Marine Mammal Commission.

⁴ Harbour porpoise strandings with signs of gillnet fishery related interactions in areas of the U.S. mid-Atlantic region not monitored by observers (A. Read, Duke University, Beaufort, N.C., pers. comm.).

Table 2. Number of fishing vessels participating in the lower Bay of Fundy gillnet fishery by season (2-wk period) and for all seasons combined from 1998-2001.

Season	1998	1999	2000	2001
June 1-15	3	1	0	1
June 16-30	4	3	1	1
July 1-15	11	8	9	12
July 16-31	15	8	10	12
Aug 1-15	16	7	10	5
Aug 16-31	14	9	8	3
Sep 1-15	12	8	8	4
Sep 16-30	9	8	4	0
All seasons	22	11	13	13

Table 3. Summary of fishing effort (trips reported) by the lower Bay of Fundy demersal gillnet fishery from 1998-2001. Numbers in parentheses represent proportion of reported trips that were observed.

1998

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
June 1-15	0	0	4 (0.00)	0	0	7 (0.00)	11 (0.00)
June 16-30	0	0	2 (0.00)	0	0	3 (0.00)	5 (0.00)
July 1-15	45 (0.22)	0	6 (0.33)	5 (0.00)	0	6 (0.17)	62 (0.21)
July 16-31	27 (0.70)	6 (0.67)	18 (0.11)	1 (0.00)	0	3 (0.33)	55 (0.47)
Aug 1-15	16 (0.38)	6 (1.00)	17 (0.18)	1 (0.00)	0	6 (0.33)	46 (0.37)
Aug 16-31	14 (0.14)	4 (0.75)	22 (0.14)	1 (0.00)	0	2 (0.00)	43 (0.19)
Sep 1-15	11 (0.00)	0	18 (0.06)	0	0	1 (0.00)	30 (0.03)
Sep 16-31	5 (0.00)	1 (0.00)	15 (0.00)	0	0	3 (0.67)	24 (0.08)
Total	118 (0.31)	17 (0.76)	102 (0.11)	8 (0.00)	0	31 (0.19)	276 (0.24)

1999

June 1-15	0	0	0	0	0	1 (0.00)	1 (0.00)
June 16-30	0	2 (0.00)	6 (0.00)	0	0	0	8 (0.00)
July 1-15	28 (0.00)	6 (1.00)	12 (0.00)	0	0	1 (0.00)	47 (0.13)
July 16-31	43 (0.00)	17 (0.59)	10 (0.00)	0	0	3 (0.00)	73 (0.14)
Aug 1-15	18 (0.00)	8 (0.25)	8 (0.00)	0	1 (0.00)	1 (1.00)	36 (0.08)
Aug 16-31	16 (0.00)	9 (0.33)	2 (0.00)	7 (0.00)	0	7 (0.26)	41 (0.12)
Sep 1-15	2 (0.00)	4 (1.00)	10 (0.00)	5 (0.00)	0	4 (0.00)	25 (0.16)
Sep 16-31	1 (0.00)	7 (0.14)	2 (0.00)	6 (0.00)	0	11 (0.00)	27 (0.04)
Total	108 (0.00)	53 (0.49)	50 (0.00)	18 (0.00)	1 (0.00)	28 (0.11)	258 (0.11)

Table 3 (cont.)

2000

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
June 1-15	0	0	0	0	0	0	0
June 16-30	0	1 (0.00)	0	0	0	0	1 (0.00)
July 1-15	12 (0.00)	2 (0.00)	7 (0.00)	0	0	4 (0.00)	25 (0.00)
July 16-31	31 (1.00)	8 (0.00)	30 (0.00)	2 (0.00)	0	3 (0.00)	74 (0.42)
Aug 1-15	44 (1.00)	2 (0.00)	3 (0.00)	0	0	5 (0.00)	54 (0.81)
Aug 16-31	37 (0.70)	16 (0.00)	14 (0.00)	0	0	2 (0.00)	69 (0.38)
Sep 1-15	10 (0.10)	4 (0.00)	11 (0.00)	0	0	3 (0.00)	28 (0.04)
Sep 16-31	0	0	0	0	0	0	0
Total	134 (0.76)	33 (0.00)	65 (0.00)	2 (0.00)	0	17 (0.00)	251 (0.41)

2001

June 1-15	0	0	2 (0.00)	0	0	0	2 (0.00)
June 16-30	0	0	4 (0.00)	0	0	0	4 (0.00)
July 1-15	44 (0.41)	5 (0.00)	14 (0.00)	4 (0.00)	0	0	67 (0.27)
July 16-31	53 (0.74)	18 (0.00)	4 (0.00)	1 (0.00)	0	3 (0.00)	79 (0.49)
Aug 1-15	36 (0.81)	6 (0.00)	0	0	0	0	42 (0.69)
Aug 16-31	35 (1.00)	0	0	0	0	0	35 (1.00)
Sep 1-15	24 (1.00)	0	0	0	0	4 (0.00)	28 (0.86)
Sep 16-31	0	0	0	0	0	0	0
Total	192 (0.76)	53 (0.00)	24 (0.00)	5 (0.00)	0	7 (0.00)	257 (0.56)

Table 4. Summary of observed porpoise catches in the lower Bay of Fundy demersal gillnet fishery from 1998-2001.

Year	Date	Area	Mesh	Porpoise	Latitude (°N)	Longitude (°W)
1998	07-Jul-98	Swallowtail	Nylon	3	44.8067	66.6596
1998	11-Jul-98	Gravelly Bulkhead	Nylon	1	-	-
1998	04-Aug-98	Swallowtail	Nylon	1	44.7904	66.6703
1999	23-Jul-99	Wolves	Mixed	1	44.9117	66.7550
1999	23-Aug-99	Wolves	Nylon	1	44.9067	66.7767
1999	14-Sep-99	Wolves	Mixed	1	44.9117	66.7450
2000	28-Jul-00	Swallowtail	Nylon	1	44.8125	66.6960
2000	01-Aug-00	Swallowtail	Nylon	1	44.8323	66.6877
2000	09-Aug-00	Swallowtail	Nylon	1	44.8208	66.7127
2000	11-Aug-00	Swallowtail	Nylon	1	44.1467	66.7198
2000	11-Aug-00	Swallowtail	Nylon	1	44.8032	66.7079
2001	10-Jul-01	Swallowtail	Barium-sulphate	1	44.8168	66.7047
2001	14-Jul-01	Swallowtail	Nylon	1	44.7983	66.6837
2001	14-Jul-01	Swallowtail	Nylon	1	44.8001	66.7032
2001	16-Jul-01	Swallowtail	Nylon	1	44.7964	66.6826
2001	16-Jul-01	Swallowtail	Nylon	1	44.7996	66.7004
2001	16-Jul-01	Swallowtail	Nylon	2	44.7992	66.6945
2001	16-Jul-01	Swallowtail	Barium-sulphate	1	44.8134	66.7121
2001	16-Jul-01	Swallowtail	Nylon	1	44.7993	66.7002
2001	16-Jul-01	Swallowtail	Barium-sulphate	1	44.8082	66.6945
2001	17-Jul-01	Swallowtail	Nylon	2	44.7978	66.6949
2001	17-Jul-01	Swallowtail	Barium-sulphate	1	44.8010	66.7164
2001	17-Jul-01	Swallowtail	Barium-sulphate	1	44.8067	66.6835
2001	18-Jul-01	Swallowtail	Nylon	1	44.7985	66.6837
2001	18-Jul-01	Swallowtail	Nylon	1	-	-
2001	20-Jul-01	Swallowtail	Barium-sulphate	1	44.8090	66.6762
Subtotal				30		

Table 4 (cont.).

Year	Date	Area	Mesh	Porpoise	Latitude (°N)	Longitude (°W)
2001	20-Jul-01	Swallowtail	Nylon	3	44.8017	66.6937
2001	20-Jul-01	Swallowtail	Barium-sulphate	1	44.8188	66.7042
2001	21-Jul-01	Swallowtail	Nylon	1	44.7967	66.6872
2001	21-Jul-01	Swallowtail	Barium-sulphate	1	44.7971	66.6821
2001	21-Jul-01	Swallowtail	Barium-sulphate	1	44.8018	66.6903
2001	30-Jul-01	Swallowtail	Nylon	1	44.7991	66.6954
2001	11-Aug-01	Swallowtail	Nylon	1	44.8368	66.6985
2001	15-Aug-01	Swallowtail	Barium-sulphate	1	44.8074	66.6986
2001	16-Aug-01	Swallowtail	Barium-sulphate	1	44.8088	66.6926
2001	18-Aug-01	Swallowtail	Nylon	2	44.7995	66.7013
2001	29-Aug-01	Swallowtail	Nylon	1	44.8362	66.7065
2001	01-Sep-01	Swallowtail	Barium-sulphate	1	44.8333	66.6983
2001	04-Sep-01	Swallowtail	Barium-sulphate	1	44.8096	66.7152
2001	05-Sep-01	Swallowtail	Nylon	1	44.8096	66.7152
2001	05-Sep-01	Swallowtail	Nylon	1	44.8029	66.6983
2001	06-Sep-01	Swallowtail	Nylon	1	44.8020	66.6977
2001	07-Sep-01	Swallowtail	Barium-sulphate	1	44.7540	66.7224
2001	08-Sep-01	Swallowtail	Barium-sulphate	1	44.7936	66.7149
2001	10-Sep-01	Swallowtail	Barium-sulphate	1	44.7908	66.6894
Total				52		

Table 5. Observed effort (trips observed) and porpoise catches of both nylon and barium-sulphate gillnet mesh nets from 1998-2001.

1998

Season	Location	Trips (#)		Catch (#)		Catch Rate (Porpoises/Trip)	
		Nylon	Barium-sulphate	Nylon	Barium-sulphate	Nylon	Barium-sulphate
July 1-15	Swallowtail	10	10	3	0	0.3000	0.0000
July 16-31	Swallowtail	14	19	0	0	0.0000	0.0000
Aug 1-15	Swallowtail	6	4	1	0	0.1667	0.0000
Aug 16-31	Swallowtail	2	2	0	0	0.1667	0.0000
July 16-31	Wolves	4	1	0	0	0.0000	0.0000
Aug 1-15	Wolves	6	5	0	0	0.0000	0.0000
Aug 16-31	Wolves	3	2	0	0	0.0000	0.0000
July 1-15	Gravelly Bulkhead	2	0	1	0	0.5000	0.0000
July 16-31	Gravelly Bulkhead	2	0	0	0	0.0000	0.0000
Aug 1-15	Gravelly Bulkhead	3	0	0	0	0.0000	0.0000
Aug 16-31	Gravelly Bulkhead	3	2	0	0	0.0000	0.0000
Sep 1-15	Gravelly Bulkhead	1	1	0	0	0.0000	0.0000
July 1-15	Head & Horns	1	1	0	0	0.0000	0.0000
July 16-31	Head & Horns	1	1	0	0	0.0000	0.0000
Aug 1-15	Head & Horns	2	0	0	0	0.0000	0.0000
July 16-31	Grand Manan Banks	2	1	0	0	0.0000	0.0000
Total		62	49	5	0		

Table 5 (cont.)

1999

Season	Location	Trips (#)			Catch (#)			Catch Rate (Porpoises/Trip)		
		Nylon	Barium-sulphate	Mixed	Nylon	Barium-sulphate	Mixed	Nylon	Barium-sulphate	Mixed
July 1-15	Wolves	6	6	4	0	0	0	0.0000	0.0000	0.0000
July 16-31	Wolves	9	0	10	0	0	1	0.0000	0.0000	0.1000
Aug 1-15	Wolves	2	0	0	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Wolves	3	0	0	1	0	0	0.3333	0.0000	0.0000
Sep 1-15	Wolves	5	0	5	0	0	1	0.0000	0.0000	0.2000
Aug 1-15	Basin	1	1	1	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Basin	2	2	1	0	0	0	0.0000	0.0000	0.0000
Aug 1-15	Digby Neck	3	3	1	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Digby Neck	11	10	7	0	0	0	0.0000	0.0000	0.0000
Totals		42	22	29	1	0	2			

Table 5 (cont.)

2000

Season	Location	Trips (#)		Catch (#)		Catch Rate (Porpoises/Trip)	
		Nylon	Barium-sulphate	Nylon	Barium-sulphate	Nylon	Barium-sulphate
July 16-31	Swallowtail	31	21	1	0	0.0323	0.0000
Aug 1-15	Swallowtail	44	44	4	0	0.0909	0.0000
Aug 16-31	Swallowtail	26	26	0	0	0.0000	0.0000
Sep 1-15	Swallowtail	1	1	0	0	0.0000	0.0000
	Total	102	92	5	0		

2001

July 1-15	Swallowtail	18	18	2	1	0.1111	0.0556
July 16-31	Swallowtail	39	38	14	8	0.3590	0.2105
Aug 1-15	Swallowtail	27	29	1	1	0.0370	0.0345
Aug 16-31	Swallowtail	35	35	3	1	0.0857	0.0286
Sep 1-15	Swallowtail	22	24	3	5	0.1364	0.2083
	Total	141	144	23	16		

Table 6. Observed effort (strings observed) and porpoise catches of both nylon and barium-sulphate gillnet mesh nets from 1998-2001.

1998

Season	Location	Trips (#)		Catch (#)		Catch Rate (Porpoises/String)	
		Nylon	Barium-sulphate	Nylon	Barium-sulphate	Nylon	Barium-sulphate
July 1-15	Swallowtail	31	12	3	0	0.0968	0.0000
July 16-31	Swallowtail	45	27	0	0	0.0000	0.0000
Aug 1-15	Swallowtail	19	5	1	0	0.0526	0.0000
Aug 16-31	Swallowtail	7	2	0	0	0.0000	0.0000
July 16-31	Wolves	19	1	0	0	0.0000	0.0000
Aug 1-15	Wolves	24	5	0	0	0.0000	0.0000
Aug 16-31	Wolves	9	2	0	0	0.0000	0.0000
July 1-15	Gravelly Bulkhead	5	0	1	0	0.2000	0.0000
July 16-31	Gravelly Bulkhead	2	0	0	0	0.0000	0.0000
Aug 1-15	Gravelly Bulkhead	3	0	0	0	0.0000	0.0000
Aug 16-31	Gravelly Bulkhead	6	2	0	0	0.0000	0.0000
Sep 1-15	Gravelly Bulkhead	3	2	0	0	0.0000	0.0000
July 1-15	Head & Horns	3	1	0	0	0.0000	0.0000
July 16-31	Head & Horns	2	1	0	0	0.0000	0.0000
Aug 1-15	Head & Horns	6	0	0	0	0.0000	0.0000
July 16-31	Grand Manan Banks	4	1	0	0	0.0000	0.0000
	Total	188	61	5	0		

Table 6 (cont).

1999

Season	Location	Trips (#)			Catch (#)			Catch Rate (Porpoises/String)		
		Nylon	Barium-sulphate	Mixed	Nylon	Barium-sulphate	Mixed	Nylon	Barium-sulphate	Mixed
July 1-15	Wolves	16	8	6	0	0	0	0.0000	0.0000	0.0000
July 16-31	Wolves	22	0	26	0	0	1	0.0000	0.0000	0.0385
Aug 1-15	Wolves	8	0	0	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Wolves	12	0	0	1	0	0	0.0833	0.0000	0.0000
Sep 1-15	Wolves	20	0	9	0	0	1	0.0000	0.0000	0.1111
Aug 1-15	Basin	2	3	2	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Basin	23	13	3	0	0	0	0.0000	0.0000	0.0000
Aug 1-15	Digby Neck	3	3	1	0	0	0	0.0000	0.0000	0.0000
Aug 16-31	Digby Neck	26	10	7	0	0	0	0.0000	0.0000	0.0000
Totals		132	37	54	1	0	2			

2000

Season	Location	Trips (#)		Catch (#)		Catch Rate (Porpoises/String)	
		Nylon	Barium-sulphate	Nylon	Barium-sulphate	Nylon	Barium-sulphate
July 16-31	Swallowtail	46	25	1	0	0.0217	0.0000
Aug 1-15	Swallowtail	84	51	4	0	0.0476	0.0000
Aug 16-31	Swallowtail	80	49	0	0	0.0000	0.0000
Sep 1-15	Swallowtail	2	2	0	0	0.0000	0.0000
Total		212	127	5	0		

2001

Season	Location	Trips (#)		Catch (#)		Catch Rate (Porpoises/String)	
		Nylon	Barium-sulphate	Nylon	Barium-sulphate	Nylon	Barium-sulphate
July 1-15	Swallowtail	49	50	2	1	0.0408	0.0200
July 16-31	Swallowtail	112	111	14	8	0.1250	0.0721
Aug 1-15	Swallowtail	75	87	1	1	0.0133	0.0115
Aug 16-31	Swallowtail	92	99	3	1	0.0326	0.0101
Sep 1-15	Swallowtail	60	68	3	5	0.0500	0.0735
Total		388	415	23	16		

Table 7. Porpoise catch rates adjusted from catch per string to catch per trip (five strings per trip) from 1998-2001.

1998			
Season	Location	Nylon Catch Rate (Porpoises/Trip)	Barium-sulphate Catch Rate (Porpoises/Trip)
July 1-15	Swallowtail	0.4839	0.0000
July 16-31	Swallowtail	0.0000	0.0000
Aug 1-15	Swallowtail	0.2632	0.0000
Aug 16-31	Swallowtail	0.0000	0.0000
July 16-31	Wolves	0.0000	0.0000
Aug 1-15	Wolves	0.0000	0.0000
Aug 16-31	Wolves	0.0000	0.0000
July 1-15	Head & Horns	0.0000	0.0000
July 16-31	Head & Horns	0.0000	0.0000
Aug 1-15	Head & Horns	0.0000	0.0000
July 1-15	Gravelly	1.0000	0.0000
July 16-31	Gravelly	0.0000	0.0000
Aug 1-15	Gravelly	0.0000	0.0000
Aug 16-31	Gravelly	0.0000	0.0000
Sep 1-15	Gravelly	0.0000	0.0000
July 16-31	Banks	0.0000	0.0000

1999				
Season	Location	Nylon Catch Rate (Porpoises/Trip)	Barium-sulphate Catch Rate (Porpoises/Trip)	Mixed Catch Rate (Porpoises/Trip)
Aug 1-15	Basin	0.0000	0.0000	0.0000
Aug 16-31	Basin	0.0000	0.0000	0.0000
Aug 1-15	Digby Neck	0.0000	0.0000	0.0000
Aug 16-31	Digby Neck	0.0000	0.0000	0.0000
July 1-15	Wolves	0.0000	0.0000	0.0000
July 16-31	Wolves	0.0000	0.0000	0.1923
Aug 1-15	Wolves	0.0000	0.0000	0.0000
Aug 16-31	Wolves	0.4167	0.0000	0.0000
Sep 1-22	Wolves	0.0000	0.0000	0.5556

Table 7 (cont.)

2000

Season	Location	Nylon Catch Rate (Porpoises/Trip)	Barium-sulphate Catch Rate (Porpoises/Trip)
July 16-31	Swallowtail	0.1087	0.0000
Aug 1-15	Swallowtail	0.2381	0.0000
Aug 16-31	Swallowtail	0.0000	0.0000
Sep 1-15	Swallowtail	0.0000	0.0000

2001

Season	Location	Nylon Catch Rate (Porpoises/Trip)	Barium-sulphate Catch Rate (Porpoises/Trip)
July 1-15	Swallowtail	0.2041	0.1000
July 16-31	Swallowtail	0.6250	0.3604
Aug 1-15	Swallowtail	0.0667	0.0575
Aug 16-31	Swallowtail	0.1630	0.0505
Sep 1-15	Swallowtail	0.2500	0.3676

Table 8. Observed catch rates (porpoises/trip) used to calculate proportional catch rates between fishing grounds.

1994

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15						
July 16-31	0.5000	0.0833		0.0000	0.0000	0.0000
Aug 1-15	0.5500	0.5000		0.5000	0.0909	0.0000
Aug 16-31	0.6111	0.6111		1.0000	0.0000	0.0000
Sep 1-15	0.2000	0.2000		0.0000	0.0000	0.0000
Sep 16-30						

1995

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15	0.3864					
July 16-31						
Aug 1-15						
Aug 16-31						
Sep 1-15	0.3636					
Sep 16-30						

1997

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15						
July 16-31						
Aug 1-15	0.2391	0.0000		0.4706		
Aug 16-31						
Sep 1-15						
Sep 16-30						

1998

Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15	0.4839		1.0000			0.0000
July 16-31	0.0000	0.0000	0.0000			0.0000
Aug 1-15	0.2632	0.0000	0.0000			0.0000
Aug 16-31	0.0000	0.0000	0.0000			
Sep 1-15			0.0000			
Sep 16-30						

Table 8 (cont.)

1999						
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15		0.0000				
July 16-31		0.0000				
Aug 1-15		0.0000				0.0000
Aug 16-31		0.4167				0.0000
Sep 1-15		0.0000				
Sep 16-30		0.0000				
2000						
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15						
July 16-31	0.1087					
Aug 1-15	0.2381					
Aug 16-31	0.0000					
Sep 1-15	0.0000					
Sep 16-30						
2001						
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other
July 1-15	0.2041					
July 16-31	0.6250					
Aug 1-15	0.0667					
Aug 16-31	0.1630					
Sep 1-15	0.2500					
Sep 16-30						

Table 9. Mean ratio of observed catch rates in each fishing area (or season) relative to Swallowtail catch rates (or Aug 1-15). N = number of seasonal pairs between a particular fishing area and Swallowtail for which the mean ratio of catch rates could be calculated.

Mean ratio relative to Swallowtail

Location	Mean Proportion of Swallowtail	N	Standard Error
Wolves	0.4849	6	0.2092
Gravelly Bulkhead	1.1810	2	1.0229
Head Harbour	1.3591	5	0.3984
Channel	0.0615	4	0.0468

Mean ratio relative to Aug 1-15 within Swallowtail

Season	Mean Proportion of Aug 1-15	N	Standard Error
July 1-15	2.7406	2	0.5378
July 16-31	2.8997	4	2.2922
Aug 15-31	0.7127	4	0.5854
Sep 1-15	1.2053	3	1.1730

Table 10. Mean catch rates (porpoises/trip) calculated from observed catch rates between Swallowtail and other fishing grounds. Numbers in bold italics indicate catch rates were calculated from observed data. Numbers not in bold italics were calculated by pro-rating standard catch rates.

1998

Season	Swallowtail	Wolves	Gravelly			
			Bulkhead	Head Harbour	Channel	Other
July 1-15	0.4839	0.2346	1.0000	0.6576	0.0298	0.0000
July 16-31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Aug 1-15	0.2632	0.0000	0.0000	0.3577	0.0162	0.0000
Aug 16-31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Sep 1-15	0.3172	0.1538	0.0000	0.4311	0.4311	0.0000

1999

Season	Swallowtail	Wolves	Gravelly			
			Bulkhead	Head Harbour	Channel	Other
July 1-15	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
July 16-31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Aug 1-15	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Aug 16-31	0.8593	0.4167	1.0148	1.1679	0.0528	0.0000
Sep 1-15	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

2000

Season	Swallowtail	Wolves	Gravelly			
			Bulkhead	Head Harbour	Channel	Other
July 1-15	0.6525	0.3164	0.7706	0.8868	0.0401	0.0000
July 16-31	0.1087	0.0527	0.1284	0.1477	0.0067	0.0000
Aug 1-15	0.2381	0.1155	0.2812	0.3236	0.0146	0.0000
Aug 16-31	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Sep 1-15	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

2001

Season	Swallowtail	Wolves	Gravelly			
			Bulkhead	Head Harbour	Channel	Other
July 1-15	0.2041	0.0990	0.2410	0.2774	0.0033	0.0000
July 16-31	0.6250	0.3031	0.7381	0.8494	0.0101	0.0000
Aug 1-15	0.0667	0.0323	0.0787	0.0906	0.0011	0.0000
Aug 16-31	0.1630	0.0791	0.1926	0.2216	0.0026	0.0000
Sep 1-15	0.2500	0.1212	0.2953	0.3398	0.0040	0.0000

Table 11. Total estimated porpoise by-catch in the lower Bay of Fundy demersal gillnet fishery from 1998-2001. Numbers in parentheses is proportion of annual estimated by-catch (small rounding errors may be present).

1998							
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
July 1-15	20 (0.526)	0	5 (0.132)	4 (0.105)	0	0	29 (0.763)
July 16-31	0	0	0	0	0	0	0
Aug 1-15	4 (0.105)	0	0	1 (0.026)	0	0	5 (0.132)
Aug 16-31	0	0	0	0	0	0	0
Sep 1-15	4 (0.105)	0	0	0	0	0	4 (0.105)
Total	28 (0.737)	0	5 (0.132)	5 (0.132)	0	0	38

1999							
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
July 1-15	0	0	0	0	0	0	0
July 16-31	0	1 (0.031)	0	0	0	0	1 (0.031)
Aug 1-15	0	0	0	0	0	0	0
Aug 16-31	14 (0.438)	4 (0.125)	3 (0.094)	9 (0.281)	0	0	30 (0.938)
Sep 1-15	0	1 (0.031)	0	0	0	0	1 (0.031)
Total	14 (0.438)	6 (0.188)	3 (0.094)	9 (0.281)	0	0	32

2000							
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
July 1-15	8 (0.286)	1 (0.036)	6 (0.214)	0	0	0	15 (0.536)
July 16-31	1 (0.036)	1 (0.036)	4 (0.143)	1 (0.036)	0	0	7 (0.250)
Aug 1-15	4 (0.143)	1 (0.036)	1 (0.036)	0	0	0	6 (0.214)
Aug 16-31	0	0	0	0	0	0	0
Sep 1-15	0	0	0	0	0	0	0
Total	13 (0.464)	3 (0.107)	11 (0.393)	1 (0.036)	0	0	28

2001							
Season	Swallowtail	Wolves	Gravelly Bulkhead	Head Harbour	Channel	Other	Total
July 1-15	9 (0.123)	1 (0.014)	4 (0.055)	2 (0.027)	0	0	16 (0.219)
July 16-31	31 (0.425)	6 (0.082)	3 (0.041)	1 (0.014)	0	0	41 (0.562)
Aug 1-15	3 (0.041)	1 (0.014)	0	0	0	0	4 (0.055)
Aug 16-31	4 (0.055)	0	0	0	0	0	4 (0.055)
Sep 1-15	8 (0.110)	0	0	0	0	0	8 (0.110)
Total	55 (0.753)	8 (0.110)	7 (0.096)	3 (0.041)	0	0	73




Fig. 1. Map of lower Bay of Fundy listing areas of traditional fishing grounds.




Fig. 2. Map of all observed porpoise mortalities in the lower Bay of Fundy demersal gillnet fishery from 1998-2001 ($n=52$, some overlap exists). White squares represent mortalities in 100% nylon-mesh nets while solid squares represent mortalities in barium-sulphate nets. Fishing areas used to partition effort for the estimation of by-catch are indicated.




Fig. 3. Total estimated porpoise by-catch in the lower Bay of Fundy demersal gillnet fishery from 1994-2001.




Fig. 4. Temporal trends in estimated porpoise by-catch in the lower Bay of Fundy demersal gillnet fishery from 1998-2001 within four fishing grounds.




Fig. 5. Temporal trends in observed and total estimated porpoise by-catch rate in the lower Bay of Fundy demersal gillnet fishery from 1998-2001.